

AD-A169 251


ASYMPTOTIC AND NUMERICAL METHODS FOR SINGULARLY
PERTURBED DIFFERENTIAL EQ. (U) RENSSELAER POLYTECHNIC
INST TROY NY DEPT OF MATHEMATICAL SCIE.
J E FLAHERTY ET AL. 19 MAY 86


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY

CHART

②

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARO 19512.17-MA	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) Asymptotic and Numerical Methods for Singularly Perturbed Differential Equations with Applications to Impact Problems		5. TYPE OF REPORT & PERIOD COVERED Final Report 10/1/82 - 1/31/86
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Joseph E. Flaherty Robert E. O'Malley		8. CONTRACT OR GRANT NUMBER(s) DAAG29-82-K-0197
9. PERFORMING ORGANIZATION NAME AND ADDRESS Rensselaer Polytechnic Institute Departments of Mathematical and Computer Science Troy, New York 12180-3590		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709 MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 5/19/86
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We developed and applied numerical methods for singularly perturbed two-point boundary value problems and time-dependent partial differential equations. DTIC FILE COPY		

AD-A169 251

86 7 1 021

FINAL REPORT

U. S. Army Research Office Contract No. DAAG29-82-K-0197

Period: 1 October 1982 through 31 January 1986

Title of Research: Asymptotic and Numerical Methods for
Singularly Perturbed Differential
Equations with Applications to Impact
Problems

Principal Investigators: Joseph E. Flaherty
Robert E. O'Malley, Jr.

Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

Approved For	
EX-100	<input checked="" type="checkbox"/>
EX-101	<input type="checkbox"/>
EX-102	<input type="checkbox"/>
EX-103	
EX-104	
EX-105	
EX-106	
EX-107	
EX-108	
EX-109	
EX-110	
EX-111	
EX-112	
EX-113	
EX-114	
EX-115	
EX-116	
EX-117	
EX-118	
EX-119	
EX-120	
EX-121	
EX-122	
EX-123	
EX-124	
EX-125	
EX-126	
EX-127	
EX-128	
EX-129	
EX-130	
EX-131	
EX-132	
EX-133	
EX-134	
EX-135	
EX-136	
EX-137	
EX-138	
EX-139	
EX-140	
EX-141	
EX-142	
EX-143	
EX-144	
EX-145	
EX-146	
EX-147	
EX-148	
EX-149	
EX-150	
EX-151	
EX-152	
EX-153	
EX-154	
EX-155	
EX-156	
EX-157	
EX-158	
EX-159	
EX-160	
EX-161	
EX-162	
EX-163	
EX-164	
EX-165	
EX-166	
EX-167	
EX-168	
EX-169	
EX-170	
EX-171	
EX-172	
EX-173	
EX-174	
EX-175	
EX-176	
EX-177	
EX-178	
EX-179	
EX-180	
EX-181	
EX-182	
EX-183	
EX-184	
EX-185	
EX-186	
EX-187	
EX-188	
EX-189	
EX-190	
EX-191	
EX-192	
EX-193	
EX-194	
EX-195	
EX-196	
EX-197	
EX-198	
EX-199	
EX-200	
EX-201	
EX-202	
EX-203	
EX-204	
EX-205	
EX-206	
EX-207	
EX-208	
EX-209	
EX-210	
EX-211	
EX-212	
EX-213	
EX-214	
EX-215	
EX-216	
EX-217	
EX-218	
EX-219	
EX-220	
EX-221	
EX-222	
EX-223	
EX-224	
EX-225	
EX-226	
EX-227	
EX-228	
EX-229	
EX-230	
EX-231	
EX-232	
EX-233	
EX-234	
EX-235	
EX-236	
EX-237	
EX-238	
EX-239	
EX-240	
EX-241	
EX-242	
EX-243	
EX-244	
EX-245	
EX-246	
EX-247	
EX-248	
EX-249	
EX-250	
EX-251	
EX-252	
EX-253	
EX-254	
EX-255	
EX-256	
EX-257	
EX-258	
EX-259	
EX-260	
EX-261	
EX-262	
EX-263	
EX-264	
EX-265	
EX-266	
EX-267	
EX-268	
EX-269	
EX-270	
EX-271	
EX-272	
EX-273	
EX-274	
EX-275	
EX-276	
EX-277	
EX-278	
EX-279	
EX-280	
EX-281	
EX-282	
EX-283	
EX-284	
EX-285	
EX-286	
EX-287	
EX-288	
EX-289	
EX-290	
EX-291	
EX-292	
EX-293	
EX-294	
EX-295	
EX-296	
EX-297	
EX-298	
EX-299	
EX-300	
EX-301	
EX-302	
EX-303	
EX-304	
EX-305	
EX-306	
EX-307	
EX-308	
EX-309	
EX-310	
EX-311	
EX-312	
EX-313	
EX-314	
EX-315	
EX-316	
EX-317	
EX-318	
EX-319	
EX-320	
EX-321	
EX-322	
EX-323	
EX-324	
EX-325	
EX-326	
EX-327	
EX-328	
EX-329	
EX-330	
EX-331	
EX-332	
EX-333	
EX-334	
EX-335	
EX-336	
EX-337	
EX-338	
EX-339	
EX-340	
EX-341	
EX-342	
EX-343	
EX-344	
EX-345	
EX-346	
EX-347	
EX-348	
EX-349	
EX-350	
EX-351	
EX-352	
EX-353	
EX-354	
EX-355	
EX-356	
EX-357	
EX-358	
EX-359	
EX-360	
EX-361	
EX-362	
EX-363	
EX-364	
EX-365	
EX-366	
EX-367	
EX-368	
EX-369	
EX-370	
EX-371	
EX-372	
EX-373	
EX-374	
EX-375	
EX-376	
EX-377	
EX-378	
EX-379	
EX-380	
EX-381	
EX-382	
EX-383	
EX-384	
EX-385	
EX-386	
EX-387	
EX-388	
EX-389	
EX-390	
EX-391	
EX-392	
EX-393	
EX-394	
EX-395	
EX-396	
EX-397	
EX-398	
EX-399	
EX-400	
EX-401	
EX-402	
EX-403	
EX-404	
EX-405	
EX-406	
EX-407	
EX-408	
EX-409	
EX-410	
EX-411	
EX-412	
EX-413	
EX-414	
EX-415	
EX-416	
EX-417	
EX-418	
EX-419	
EX-420	
EX-421	
EX-422	
EX-423	
EX-424	
EX-425	
EX-426	
EX-427	
EX-428	
EX-429	
EX-430	
EX-431	
EX-432	
EX-433	
EX-434	
EX-435	
EX-436	
EX-437	
EX-438	
EX-439	
EX-440	
EX-441	
EX-442	
EX-443	
EX-444	
EX-445	
EX-446	
EX-447	
EX-448	
EX-449	
EX-450	
EX-451	
EX-452	
EX-453	
EX-454	
EX-455	
EX-456	
EX-457	
EX-458	
EX-459	
EX-460	
EX-461	
EX-462	
EX-463	
EX-464	
EX-465	
EX-466	
EX-467	
EX-468	
EX-469	
EX-470	
EX-471	
EX-472	
EX-473	
EX-474	
EX-475	
EX-476	
EX-477	
EX-478	
EX-479	
EX-480	
EX-481	
EX-482	
EX-483	
EX-484	
EX-485	
EX-486	
EX-487	
EX-488	
EX-489	
EX-490	
EX-491	
EX-492	
EX-493	
EX-494	
EX-495	
EX-496	
EX-497	
EX-498	
EX-499	
EX-500	
EX-501	
EX-502	
EX-503	
EX-504	
EX-505	
EX-506	
EX-507	
EX-508	
EX-509	
EX-510	
EX-511	
EX-512	
EX-513	
EX-514	
EX-515	
EX-516	
EX-517	
EX-518	
EX-519	
EX-520	
EX-521	
EX-522	
EX-523	
EX-524	
EX-525	
EX-526	
EX-527	
EX-528	
EX-529	
EX-530	
EX-531	
EX-532	
EX-533	
EX-534	
EX-535	
EX-536	
EX-537	
EX-538	
EX-539	
EX-540	
EX-541	
EX-542	
EX-543	
EX-544	
EX-545	
EX-546	
EX-547	
EX-548	
EX-549	
EX-550	
EX-551	
EX-552	
EX-553	
EX-554	
EX-555	
EX-556	
EX-557	
EX-558	
EX-559	
EX-560	
EX-561	
EX-562	
EX-563	
EX-564	
EX-565	
EX-566	
EX-567	
EX-568	
EX-569	
EX-570	
EX-571	
EX-572	
EX-573	
EX-574	
EX-575	
EX-576	
EX-577	
EX-578	
EX-579	
EX-580	
EX-581	
EX-582	
EX-583	
EX-584	
EX-585	
EX-586	
EX-587	
EX-588	
EX-589	
EX-590	
EX-591	
EX-592	
EX-593	
EX-594	
EX-595	
EX-596	
EX-597	
EX-598	
EX-599	
EX-600	
EX-601	
EX-602	
EX-603	
EX-604	
EX-605	
EX-606	
EX-607	
EX-608	
EX-609	
EX-610	
EX-611	
EX-612	
EX-613	
EX-614	
EX-615	
EX-616	
EX-617	
EX-618	
EX-619	
EX-620	
EX-621	
EX-622	
EX-623	
EX-624	
EX-625	
EX-626	
EX-627	
EX-628	
EX-629	
EX-630	
EX-631	
EX-632	
EX-633	
EX-634	
EX-635	
EX-636	
EX-637	
EX-638	
EX-639	
EX-640	
EX-641	
EX-642	
EX-643	
EX-644	
EX-645	
EX-646	
EX-647	
EX-648	
EX-649	
EX-650	
EX-651	
EX-652	
EX-653	
EX-654	
EX-655	
EX-656	
EX-657	
EX-658	
EX-659	
EX-660	
EX-661	
EX-662	
EX-663	
EX-664	
EX-665	
EX-666	

1. Research on Numerical Methods for Singularly Perturbed Differential Equations

During the period of this contract, we developed and applied numerical methods for singularly perturbed two-point boundary value problems and time-dependent partial differential equations. Our progress and publications in these areas are described below.

1.1. Two Point Boundary Value Problems

Our research during this period has largely centered on analyzing the asymptotic behavior of solutions to scalar problems of the form

$$\epsilon y'' + f(y)y' + g(y) = 0, 0 \leq x \leq \epsilon$$

with $y(0) = A$ and $y(\epsilon) = B$ prescribed. Such problems and their generalizations model a variety of nonlinear oscillators and other physical systems involving narrow boundary and interior layers where solutions change rapidly. A most fascinating example involves $f(y) = y(1-y^2)$ and $g(y) = -y$. The unique solution differs in twenty-two different regions of the A-B plane, having up to four different limiting solutions in each (depending on the interval length ϵ). Away from regions of nonuniform convergence, the limiting solution $Y_0(x)$ will satisfy one of the five limits: $(1 - Y_0^2) \dot{Y}_0 = 1, Y_0(0) = A$; $(1 - Y_0^2) \dot{Y}_0 = 1, Y_0(0) = -1$; $(1 - Y_0^2) \dot{Y}_0 = 1, Y_0(\epsilon) = B$; $(1 - Y_0^2) \dot{Y}_0 = 1, Y_0(\epsilon) = 1$; or $Y_0(x) = 0$.

This behavior and the location of jump points can be explained by analyzing trajectories for the corresponding fast-slow system in the y-z phase plane, where z is the Lienard variable $z = \epsilon \dot{y} + F(y)$, $F(y) = \int_0^y f(s) ds$. More complicated behavior will be obtained for systems with more rest points and when the "characteristic curve" $z = F(y)$ has less symmetry and more extreme points. Our results may be typical of nonlinear problems which arise as reaction-diffusion systems or as other problems with slow/fast dynamics. They would be nearly impossible to comprehend and discover computationally, without some prior knowledge of possible analytical behavior. This work is being carried out with graduate student John Allen. It has benefited from graphical ODE routines produced by G. Odell. Further computational efforts will benefit from use of adaptive numerical methods, as have been developed by Flaherty, Brown and Lorenz, and others, and, ultimately, by devising domain decomposition techniques related to identification of asymptotic scales.

Vasil'eva and Butuzov, and Flaherty and O'Malley have independently developed analyses for singular-singular perturbation problems. These problems occur when the limiting equation has a family of solutions. They arise in

applications ranging from chemical kinetics to circuit theory. Recently, proposed simulation methods for rapidly oscillating controls made clear the need for a more comprehensive theory. This is the focus of thesis work by Gu Zhong-Mei and of a collaboration with Dr. Nikolai Nefedov, a visiting faculty member from Moscow State University. At this time, it appears that the Flaherty-O'Malley procedure (which provides a differential equation for the limiting solution) is most practical. However, more practical means of decoupling fast and slow parts of solutions and of examining appropriate stability manifolds for stretched systems are both very much needed. Results obtained will form a basis for regularization methods to solve differential-algebraic systems, as well as the original singular problems.

Other efforts by Professor O'Malley include continuing interactions with G. Soderlind of Stockholm's Royal Institute of Technology, R. Mattheij of the Catholic University of Nijmegen, and with colleagues J. Cole, J. Flaherty, H. Kurland, and M. Slemrod, among others. A fairly extensive set of class notes on singular perturbation methods was written last semester. It may soon form the basis of a research level monograph or graduate textbook.

1.2 Initial Boundary Value Problems for Partial Differential Equations

Professor Flaherty and several colleagues and graduate students have been conducting research on adaptive finite difference and finite element methods for partial differential equations.

Drew and Flaherty [2]¹ created a model for shear band instabilities that involves the rapid shearing of a slab of a visco-elastic material with small viscosity. They used a moving mesh finite element code to solve their model and illustrated several possible mechanisms for shear band formations. In a somewhat related study, Slemrod and Flaherty [12] considered appropriate finite difference schemes for phase transitions involving van der Waals fluids.

Moving mesh methods were studied and the stability of several schemes that are based on equidistributing a local error indicator was characterized. Our findings, reported in [11], show that many popular mesh moving strategies can be unstable in certain situations.

Moving grid methods are effective at reducing dispersive errors in the vicinity of wave fronts; however, no method that uses a fixed number of computational cells can be used to compute a solution to a prescribed level of accuracy. For this reason, we have been studying and developing techniques that perform explicit error estimations

¹ See list of publications and manuscripts in Section 3

that are used to add and delete elements as the temporal integration progresses. In this spirit, Adjrid and Flaherty [13] developed a finite element method of lines where the mesh "lines" are adaptively moved to minimize artificial diffusion, but where elements are added and deleted during the integration. Piecewise linear finite elements are used to calculate the numerical solution and quadratic elements are used to estimate its error. Mesh motion is accomplished using one of the stable schemes studied in [11]. Ordinary differential equations are obtained for the finite element solution, the error estimate, and the mesh coordinates and these may be solved using existing software for stiff systems. We have applied this procedure to several problems, including some difficult combustion studies, and found some very encouraging results.

The elementary global refinement strategy that was used in [13] could be inefficient in many situations. Additionally, the mesh moving algorithm of [13] required users to specify a parameter that controlled relaxation times to an "ideal" mesh that equidistributes local error estimates. We have remedied both of these deficiencies and are now using an efficient local refinement strategy and an adaptive procedure for selecting the mesh relaxation parameter. The details of these algorithms and the results of several computational experiments have been incorporated into a manuscript which has been accepted for publication in *Comput. Meths. in Appl. Mech. Engr.* [16]. We have also extended our solution, error estimation, and local refinement procedures to two-dimensional parabolic problems. Details of the method, which uses dynamic tree structures to store data, and the results of several academic and practical examples were reported in Slimane Adjrid's Ph.D. dissertation [8].

The approach of Adjrid and Flaherty [13] couples mesh motion and error estimation to the solution. This produces a method with great numerical stability; however, there are many instances where this extra effort is not necessary. Thus, we are considering adaptive local refinement techniques that use rectangular and trapezoidal space-time elements (cf. Flaherty and Moore [3,6] and Bieterman et al. [14]). These methods discretize and solve a problem for one time step using a finite element-Galerkin procedure. At the end of the time step, the discretization error is estimated, finer subgrids of space-time elements are added to regions of high error, and the problem is recursively solved again on these regions. The process terminates and the integration continues to the next time step when the estimated error on each grid is less than a prescribed tolerance.

There is more to be done with our space-time local refinement codes. Flaherty and graduate students M. J. Coyle and P. K. Moore are concentrating on developing hierarchic error estimation techniques that are similar to those of Adjrid and Flaherty [13,16] for the method of lines. The essential difficulties are (i) the absence

of any superconvergence in time and (ii) the appropriate initial and boundary conditions to apply at coarse-fine mesh interfaces. In addition, Jackson and Flaherty [17] have extended the finite element basis to include discontinuous trial functions. They used these in conjunction with a moving mesh scheme of Harten and Hyman² to develop a method that is extremely well suited to the solution of hyperbolic systems with shocks. The method captures shocks and contact surfaces as true discontinuities without any artificial diffusion or oscillations.

We have developed an adaptive finite difference method for two-dimensional systems of conservation laws. Preliminary work on a technique that used MacCormack's method on a moving grid of quadrilateral elements was reported by Arney and Flaherty [5,15]. This mesh moving technique used a clustering algorithm of Berger³ to locate and group regions of high error into rectangles that isolate spatially distinct phenomena. An algebraic mesh moving function was then used to move and align the mesh with the regions of high error. This procedure is very efficient and, unlike many other two-dimensional mesh moving techniques, it has no problem dependent parameters.

Major D. C. Arney, a professor at the U.S. Military Academy, completed his Ph.D. studies in 1985 under Flaherty's direction. He has combined the technique of [5,15] with a recursive local refinement strategy and used it to create an adaptive finite difference code for nonlinear two-dimensional hyperbolic systems [9]. The code has been applied to several difficult compressible flow problems that contain shocks, contact surfaces, and expansions. It appears to be able to utilize the best features of mesh moving and refinement.

2. Recent Interactions

Professors Flaherty and O'Malley and graduate students lectured and/or visited the following conferences and organizations during 1985.

J. E. Flaherty visited Los Alamos National Laboratory, Los Alamos, 20-21 June, 1985. He held technical discussions with Drs. J. M. Hyman, T. Manteufel, J. Dendy, B. Swartz, and B. Wendroff and lectured on "Moving Finite Difference and Finite Element Methods with Local Refinement for Time Dependent Partial Differential Equations."

² A. Harten and J. M. Hyman, "Self Adjusting Grid Methods for One-Dimensional Hyperbolic Conservation Laws," *J. Comp. Phys.*, Vol. 50, 269-325, 1983

³ M. J. Berger, "Adaptive Mesh Refinement for Hyperbolic Partial Differential Equations," Report No. STAN-CS-82-924, Department of Computer Science, Stanford University, 1982.

J. E. Flaherty attended the ASME/ASCE Applied Mechanics Conference, Albuquerque, 24-26 June, 1985. He presented an invited lecture on "Local Refinement and Moving Finite Element Methods for Parabolic Partial Differential Equations" as part of the Symposium on Recent Advances in Computational Mechanics. He also had technical discussions with Profs. R. Ewing of Wyoming University, J. T. Oden of the University of Texas, and R. D. Russell of Simon Fraser University.

R. E. O'Malley attended the AMS Symposium on Combustion Theory at Cornell University for a week in July. Especially valuable correspondence concerning singular models for chemical kinetics resulted with Professor S.-H. Lam of Princeton University.

R. E. O'Malley visited Los Alamos National Laboratory and its Center for Non-linear Studies for a week in August. He had technical discussions about singular perturbations with B. Nicolaenko, P. Fife, D. Brown, D. Cohen, J. M. Hyman, D. McLaughlin, and J. D. Murray, among others.

J. E. Flaherty co-organized (with D. A. Drew) the Conference on Mathematics Applied to Fluid Mechanics and Stability which was dedicated in memory of Professor R. C. Di Prima and held at RPI, 9-11 September, 1985.

J. E. Flaherty visited the IBM T. J. Watson Research Center, Yorktown Heights, 9 October, 1985 and lectured on "Adaptive Finite Difference and Finite Element Methods for Time Dependent Partial Differential Equations." He also had technical discussions with Drs. W. Liniger and F. Odeh.

J. E. Flaherty visited the Courant Institute of Mathematical Sciences, New York, 18 October, 1985. He had technical discussions with Profs. M. Berger and P. Collela and lectured on "Adaptive Finite Difference and Finite Element Methods for Time Dependent Partial Differential Equations."

S. Adjerid and J. E. Flaherty attended the SIAM Fall meeting, Arizona State University, Tempe, 28-30 October, 1985. They had technical discussions with several individuals, including Dr. R. C. Y. Chin of Lawrence Livermore Laboratories, Prof. R. Bank of the University of California at San Diego, and Prof. R. D. Russell of Simon Fraser University.

Professor O'Malley also attended the SIAM Fall meeting in Tempe. This provided an opportunity for discussions about numerical methods for two-point stiff problems with G. Dahlquist, R. Russell, C. Ringhofer, and C. Schmeisser, among others. In October, he gave a seminar on the subject at the University of Montreal and a series of three lectures at the University of New Hampshire.

S. Adjerid, J. E. Flaherty, and P. K. Moore attended the SIAM Conference on Parallel Processing for Scientific Computing, Norfolk, 18-21 November, 1985. They had technical discussions with several individuals including Maj. D. C. Arney of the United States Military Academy, Drs. R. C. Y. Chin and L. Petzold of Lawrence Livermore Laboratories, Dr. S. F. Davis of the Naval Surface Weapons Center, and Prof. J. Van Rosendale of the University of Utah.

S. Adjerid, J. E. Flaherty, and P. K. Moore attended a day-long workshop on adaptive methods for partial differential equations at the University of Maryland on 24 January 1986. Each of them presented a lecture on adaptive methods. Also present at the workshop were I. Babuska, J. Osborne, and E. Rank of the University of Maryland; P. Baehman and M. Shephard of RPI; J. Thomas of AFOSR; S. F. Davis and W. Szymczak of the Naval Surface Weapons Center; and other students and faculty.

Professor O'Malley is chairman of the 1986 SIAM National Meeting. Its program will feature many sessions relevant to this contract. He is also organizing a SIAM Conference on Numerical Methods for Singularly Perturbed Problems.

Effective January 1, 1986, Professor O'Malley is SIAM's Vice President for Publications. He has also joined the editorial board of the IMA Journal on Applied Mathematics.

Drs. O'Malley and Nefedov visited California in February to discuss singular perturbation problems with Professors Cohen, Lorenz, Lagerstrom, and Kreiss at Caltech; Professors Harris, Sibuya, and Sacker at the University of Southern California; and Drs. Howes and Chin at the Lawrence Livermore Laboratories.

3. List of Publications and Manuscripts

Publications:

1. R. E. O'Malley, Jr. and J. E. Flaherty, "On the Numerical Solution of Singularly-Perturbed Vector Boundary Value Problems," in *Trans. Tenth IMACS World Cong. on Syst. Simul. and Sci. Comput.*, R. Stepelman, M. Carver, R. Peskin, W. F. Ames, R. Vichnevetsky (Eds.), Vol. 1, North-Holland, New York, pp. 105-112, 1983.
2. D. A. Drew and J. E. Flaherty, "Adaptive Finite Element Methods and the Numerical Solution of Shear Band Problems," in *Phase Transitions and Material Instabilities in Solids*, M. Gurtin, Ed., Academic Press, New York, pp. 37-60, 1984.

3. J. E. Flaherty and P. K. Moore, "An Adaptive Local Refinement Finite Element Method for Parabolic Partial Differential Equations," in *Proc. Int. Conf. on Accuracy Estimates and Adaptive Refinements in Finite Element Computations*, Int. Assn. Compu. Mech., Vol. 2, Lisbon, pp. 139-152, 1984.
4. J. E. Flaherty, M. L. Frankel, V. Roytburd, G. I. Sivaskinsky, "Numerical Study of Quenching of Inward Propagating Spherical Flames," in *Combust. Sci. Tech.*, Vol. 43, pp. 245-257, 1985.
5. D. C. Arney and J. E. Flaherty, "A Mesh Moving Technique for Time Dependent Partial Differential Equations in Two-Space Dimensions," in *Trans. Second Army Conf. on Appl. Maths. and Comput.*, U. S. Army Research Office Report No. ARO 85-1, U. S. Army Research Office, Research Triangle Park, pp. 611-634, 1985.
6. J. E. Flaherty and P. K. Moore, "A Local Refinement Finite Element Method for Time Dependent Partial Differential Equations," in *Trans. Second Army Conf. on Appl. Maths. and Comput.*, U. S. Army Research Office Report No. ARO 85-1, U. S. Army Research Office, Research Triangle Park, pp. 585-596, 1985.
7. R. E. O'Malley, Jr., "On the Simultaneous Use of Asymptotic and Numerical Methods to Solve Nonlinear Two-point Problems with Boundary and Interior Layers," in *Numerical Boundary Value ODEs*, U. M. Ascher and R. D. Russell (Eds.), Birkhauser, Boston, pp. 149-172, 1985.
8. S. Adjerid, "Adaptive Finite Element Methods for Time Dependent Partial Differential Equations," Ph.D. Dissertation, Department of Mathematical Sciences, Rensselaer Polytechnic Institute, Troy, 1985.
9. D. C. Arney, "An Adaptive Mesh Algorithm for Solving Systems of Time Dependent Partial Differential Equations," Ph.D. Dissertation, Department of Mathematical Sciences, Rensselaer Polytechnic Institute, Troy, 1985.
10. R. E. O'Malley, Jr., and R. M. M. Mattheij, "Decoupling and Boundary Value Problems for Two-Time-Scale Systems," *Stiff Computation*, R. C. Aiken (Ed.), Oxford University Press, pp. 104-111, 1985.
11. J. M. Coyle, J. E. Flaherty and R. Ludwig, "On the Stability of Mesh Equidistribution Strategies for Time Dependent Partial Differential Equations," *J. Comp. Phys.*, Vol. 62, pp. 26-39, 1986.

In Press:

12. M. Slemrod and J. E. Flaherty. "Numerical Integration of a Riemann Problem for a van der Waals Fluid." *Res. Mechanica*, accepted for publication, 1984.
13. S. Adjerdid and J. E. Flaherty. "A Moving Finite Element Method with Error Estimation and Refinement for One-Dimensional Time Dependent Partial Differential Equations." *SIAM J. Numer. Anal.*, accepted for publication, 1985.
14. M. Bieterman, J. E. Flaherty, and P. K. Moore. "Adaptive Refinement Methods for Nonlinear Parabolic Partial Differential Equations." I. Babuska, O. C. Zienkiewicz, E. Arantes e Oliveira, J. R. Gago, and K. Morgan, (Eds.), *Accuracy Estimates and Adaptivity for Finite Elements*, John Wiley, London, accepted for publication, 1984.
15. D. C. Arney and J. E. Flaherty. "A Two-Dimensional Mesh Moving Technique for Time Dependent Partial Differential Equations." *J. Comp. Phys.*, accepted for publication, 1986.
16. S. Adjerdid and J. E. Flaherty. "A Moving Mesh Finite Element Method with Local Refinement for Parabolic Partial Differential Equations." *Computer Methods in Applied Mechanics and Engineering*, accepted for publication, 1985.
17. J. E. Flaherty and T. L. Jackson. "A Discontinuous Finite Element Method for Hyperbolic Systems of Conservation Laws." submitted to *SIAM J. Sci. Stat. Comput.*, September 1985.
18. "Singular Perturbation in Systems and Control." *IEEE Press*, 1986. (P. V. Kokotovic and H. K. Khalil, (Eds.), will reprint articles by Kokotovic, Sannuti, and O'Malley (Automatica 1976), O'Malley (SIAM J. Control, 1972), O'Malley (J. Math. Anal. Appl., 1974), and O'Malley and Jameson (IEEE Trans. Automatic Control, 1977).
19. R. E. O'Malley, Jr., "On Nonlinear Singularly Perturbed Initial Value Problems," submitted to *SIAM Review*, July 1985.

ON THE NUMERICAL SOLUTION OF SINGULARLY-PERTURBED VECTOR BOUNDARY VALUE PROBLEMS

Robert E. O'Malley, Jr.

and

Joseph E. Flaherty

Department of Mathematical Sciences

Rensselaer Polytechnic Institute

Troy, New York 12180 USA

ABSTRACT

Numerical procedures are developed for constructing asymptotic solutions of certain nonlinear singularly-perturbed vector two-point boundary value problems having boundary layers at one or both end points. The asymptotic approximations are generated numerically and can either be used as is or to furnish a two-point boundary value code (e.g., COLSYS) with an initial approximation and a nonuniform computational mesh. The procedures are applied to a model problem that indicates the possibility of multiple solutions and problems involving the deformation of a thin nonlinear elastic beam resting on a nonlinear elastic foundation.

ADAPTIVE FINITE ELEMENT METHODS AND THE NUMERICAL SOLUTION OF SHEAR BAND PROBLEMS

Donald A. Drew and Joseph E. Flaherty
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

Shear bands are localized regions of very high shear strain which arise as a result of high rates of loading. They occur in metal forming and cutting processes and in impact and penetration problems. In this paper, we describe a model for the formation of shear bands in simple shear that involves the description of irreversible mechanical shear and the resulting heat release. The location of a shear band is unknown in advance, and the evolution results in large gradients of displacement, velocity, and temperature. Shear band formation, therefore, offers an interesting and physically important application of a code able to resolve small-scale transient structures. In this paper, we use an adaptive finite element code to solve several problems involving shear band formation. The code automatically locates regions with large gradients and adaptively concentrates finite elements there in order to minimize approximately the development of shear bands under many circumstances and indicate some possible mechanisms for their formation.

**AN ADAPTIVE LOCAL REFINEMENT
FINITE ELEMENT METHOD FOR PARABOLIC
PARTIAL DIFFERENTIAL EQUATIONS**

Joseph E. Flaherty
Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180

and

Peter K. Moore
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

We discuss an adaptive finite element method for solving initial-boundary value problems for vector systems of partial differential equations in one space dimension and time. The method uses piecewise bilinear rectangular space-time finite elements. For each time step, the grid is automatically refined in regions where the local discretization error is estimated as being larger than a prescribed tolerance. We discuss several aspects of our algorithm, including the tree structure that is used to represent the finite element solution and grids, an error estimation technique, and initial and boundary conditions at coarse-fine mesh interfaces. We also present the results of several computational examples and experiments.

NUMERICAL STUDY OF QUENCHING OF INWARD PROPAGATING SPHERICAL FLAMES

J. E. Flaherty

Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180

and

M. L. Frankel, V. Roytburd

Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

and

G. I. Sivashinsky

Department of Mathematical Sciences
Tel-Aviv University
Tel-Aviv 69978, Israel

ABSTRACT

The phenomenon of quenching of cylindrical convergent flames is considered in the framework of constant density approximation. The parameters dependence of the effect is studied numerically using an adaptive finite element method. The numerical results are in a good agreement with predictions of theoretical analysis from (Frankel and Sivashinsky, 1984).

**A MESH MOVING TECHNIQUE
FOR TIME DEPENDENT PARTIAL DIFFERENTIAL EQUATIONS
IN TWO SPACE DIMENSIONS**

David C. Arney
Department of Mathematics
United States Military Academy
West Point, NY 10996
and
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, NY 12180
and
Joseph E. Flaherty
Applied Mathematics and Mechanics Section
Benet Weapons Laboratory
Watervliet, NY 12189
and
Department of Computer Science
Rensselaer Polytechnic Institute
Troy, NY 12180

ABSTRACT

We discuss an adaptive mesh moving technique that is used with a finite difference or finite element scheme to solve initial-boundary value problems for vector systems of partial differential equations in two space dimensions and time. The mesh moving technique is based on an algebraic node movement function determined from the *propagation of significant error regions*. The algorithm is designed to be flexible, so that it can be used with many existing finite difference or finite element methods. To test the algorithm, we implemented the mesh mover in a system code along with an initial mesh generator and a MacCormack finite volume integrator to solve hyperbolic vector systems. Results are presented for several computational examples. The moving mesh reduces dispersion errors near shocks and wave fronts and thereby can reduce the grid requirements necessary to compute accurate solutions while increasing computational efficiency.

A LOCAL REFINEMENT FINITE ELEMENT METHOD FOR TIME DEPENDENT PARTIAL DIFFERENTIAL EQUATIONS

Joseph E. Flaherty

Department of Computer Science

Rensselaer Polytechnic Institute

Troy, N.Y. 12180

and

U. S. Army Armament Research and Development Command

Large Calibre Weapon Systems Laboratory

Benet Weapons Laboratory

Watervliet, NY 12189

and

Peter K. Moore

Department of Mathematical Sciences

Rensselaer Polytechnic Institute

Troy, NY 12180

ABSTRACT

We discuss an adaptive finite element method for solving initial-boundary value problems for vector systems of partial differential equations in one space dimension and time. The method uses piecewise bilinear rectangular space-time finite elements. For each time step, the grid is automatically refined in regions where the local discretization error is estimated as being larger than a prescribed tolerance. We discuss several aspects of our algorithm, including the tree structure that is used to represent the finite element solution and grids, an error estimation technique, and initial and boundary conditions at coarse-fine mesh interfaces. We also present the results of several computational examples and experiments.

ON THE SIMULTANEOUS USE OF ASYMPTOTIC
AND NUMERICAL METHODS TO SOLVE NONLINEAR
TWO-POINT PROBLEMS WITH BOUNDARY AND INTERIOR LAYERS

R. E. O'Malley, Jr.
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

The purpose of this paper is to provide a survey concerning boundary value problems for certain systems of nonlinear singularly perturbed ordinary differential equations. The aim is to emphasize important and difficult open problems needing more study, in terms of both mathematical and numerical analysis and computational experiments. Some currently tractable problems are discussed in detail.

ADAPTIVE FINITE ELEMENT METHODS FOR TIME DEPENDENT PARTIAL DIFFERENTIAL EQUATIONS

Slimane Adjerd
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

We discuss adaptive finite element methods for solving initial-boundary problems for vector systems of partial differential equations in one and two space dimensions.

One dimensional systems are discretized using piecewise linear finite element approximations in space and a backward difference code for stiff ordinary differential systems in time. A spatial error estimate is calculated using piecewise quadratic approximations that use the superconvergence property of parabolic systems to gain computational efficiency. This error estimate is used to move and refine the finite element mesh in order to equidistribute a measure of the total spatial error and to satisfy a prescribed error tolerance. Ordinary differential equations for the spatial error estimate and the mesh motion are integrated in time using the same backward difference software that is used to determine the numerical solution of the partial differential system. The one-dimensional algorithm combines mesh moving with local refinement in a relatively efficient manner and attempts to eliminate problem-dependent numerical parameters. A variety of examples that motivate our mesh moving strategy and illustrate the performance of our algorithms are presented.

Two-dimensional systems are discretized using piecewise bilinear finite element approximations in space and the same backward difference software that is used for one-dimensional systems in time. A spatial error estimate is calculated using piecewise cubic approximations that use the superconvergence property of parabolic systems. This error estimate is used to locally refine a stationary finite element mesh in order to satisfy a prescribed spatial error tolerance.

Several linear and nonlinear examples are presented to illustrate the effectiveness of our error estimation technique and the performance of our adaptive algorithms.

AN ADAPTIVE MESH ALGORITHM FOR SOLVING SYSTEMS OF TIME DEPENDENT PARTIAL DIFFERENTIAL EQUATIONS

David C. Arney
Department of Mathematics
United States Military Academy
West Point, NY 10996
and
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

We discuss an adaptive mesh algorithm that can be used with a finite difference or finite element scheme to solve initial-boundary value problems for vector systems of time dependent partial differential equations in two space dimensions. Our algorithm combines the adaptive techniques of mesh moving, static rezoning, and local mesh refinement. The nodes of a coarse mesh of quadrilateral cells are moved by a simple algebraic node movement function, determined from the geometry and propagation of regions having statically significant discretization error or mesh movement indicators. The local mesh refinement method recursively divides cells of the moving coarse mesh within clustered regions that contain nodes with large error until a user prescribed error tolerance is satisfied. These finer grids are properly nested within the moving coarse mesh to provide for simpler data structures and interface conditions between the fine and coarse meshes.

Our procedure is designed to be flexible, so that it can be used with many existing finite difference and finite element schemes and with different error estimation procedures. To test our adaptive mesh algorithm, we implemented it in a system code with an initial mesh generator, a MacCormack finite difference scheme for hyperbolic vector systems of conservation laws, and a Richardson extrapolation based error estimation. Results are presented for several computational examples.

The moving mesh technique reduces dispersive errors near shocks and wave fronts. Therefore, it reduces the grid requirements necessary to compute accurate solutions and thus increases computational efficiency. The local mesh refinement provides smaller mesh spacings and time steps in regions where the problem is difficult to solve, thus providing increased accuracy and enabling error tolerances to be achieved.

DECOUPLING AND BOUNDARY VALUE
PROBLEMS FOR TWO-TIME-SCALE SYSTEMS

R. E. O'Malley, Jr.
Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

and

R. M. M. Mattheij
Catholic University
Nijmegen, The Netherlands

ABSTRACT

This paper presents an analytical technique for decoupling slow and fast dynamics for linear systems of ordinary differential equations. The method, based on Riccati transformations, is of substantial computational value for solving stiff two-point boundary value problems.

ON THE STABILITY OF MESH EQUIDISTRIBUTION STRATEGIES FOR TIME DEPENDENT PARTIAL DIFFERENTIAL EQUATIONS

J. Michael Coyle, Joseph E. Flaherty and Raymond Ludwig

Department of Mathematical Sciences

Rensselaer Polytechnic Institute

Troy, New York 12180 USA

ABSTRACT

We study the stability of several mesh equidistribution schemes for time dependent partial differential equations in one space dimension. The schemes move a finite difference or finite element mesh so that a given quantity is uniform over the domain. We consider mesh moving methods that are based on solving a system of ordinary differential equations for the mesh velocities and show that many of these methods are unstable with respect to an equidistributing mesh when the partial differential system is dissipative. Using linear perturbation techniques, we are able to develop simple criteria for determining the stability of a particular method and show how to construct stable differential systems for the mesh velocities. Several examples illustrating stable and unstable mesh motions are presented.

NUMERICAL INTEGRATION OF A RIEMANN PROBLEM FOR A VAN DER WAALS FLUID

M. Slemrod

and

J. E. Flaherty

Department of Mathematical Sciences

Rensselaer Polytechnic Institute

Troy, New York 12180

ABSTRACT

In two recent papers, Slemrod has suggested that the well known Lax-Friedrichs finite difference method may provide a natural method for the numerical integration of initial value problems with an anomalous equation of state, e.g., a van der Waals fluid. In this note we review these ideas and present the results of a numerical experiment which attempts to simulate the dynamics of a van der Waals like fluid.

**A MOVING FINITE ELEMENT METHOD WITH ERROR ESTIMATION
AND REFINEMENT FOR ONE-DIMENSIONAL TIME DEPENDENT
PARTIAL DIFFERENTIAL EQUATIONS**

Slimane Adjerd

Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

and

Joseph E. Flaherty
Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180

Dedicated in Memory of Richard C. DiPrima

ABSTRACT

We discuss a moving finite element method for solving vector systems of time dependent partial differential equations in one space dimension. The mesh is moved so as to equidistribute the spatial component of the discretization error in H^1 . We present a method of estimating this error by using p-hierarchic finite elements. The error estimate is also used in an adaptive mesh refinement procedure to give an algorithm that combines mesh movement and refinement. We discretize the partial differential equations in space using a Galerkin procedure with piecewise linear elements to approximate the solution and quadratic elements to estimate the error. A system of ordinary differential equations for mesh velocities are used to control element motions. We use existing software for stiff ordinary differential equations for the temporal integration of the solution, the error estimate, and the mesh motion. Computational results using a code based on our method are presented for several examples.

ADAPTIVE REFINEMENT METHODS FOR NONLINEAR PARABOLIC PARTIAL DIFFERENTIAL EQUATIONS

Michael Bieterman

Boeing Computer Services
Engineering Technology Applications
Tukwila, Washington 98118 USA

and

Joseph E. Flaherty

Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180 USA

and

Peter K. Moore

Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180 USA

ABSTRACT

We consider two adaptive finite element techniques for parabolic partial differential equations (PDEs) that are based on using error estimates to control mesh refinement. One technique is a method of lines (MOL) approach that uses a Galerkin method to discretize the PDEs in space and implicit multi-step integration in time. Spatial elements are added and deleted in regions of high and low error and are all advanced with the same sequence of varying time steps. The second technique is a local refinement method (LRM) that uses Galerkin approximations in both space and time. Fine grids of space-time elements are added to coarser grids and the problem is recursively solved in regions of high error.

A TWO-DIMENSIONAL MESH MOVING TECHNIQUE FOR TIME DEPENDENT PARTIAL DIFFERENTIAL EQUATIONS

David C. Arney

Department of Mathematics
United States Military Academy
West Point, NY 10996

and

Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

and

Joseph E. Flaherty
Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

We discuss an adaptive mesh moving technique that can be used with a finite element difference or finite element scheme to solve initial-boundary value problems for vector systems of partial differential equations in two space dimensions and time. The mesh moving technique is based on an algebraic node movement function determined from the geometry and propagation of regions having significant discretization error indicators. Our procedure is designed to be flexible, so that it can be used with many existing finite difference and finite element methods. To test the mesh moving algorithm, we implemented it in a system code with an initial mesh generator and a MacCormack finite difference scheme on quadrilateral cells for hyperbolic vector systems of conservation laws. Results are presented for several computational examples. The moving mesh scheme reduces dispersive errors near shocks and wave fronts and thereby reduces the grid requirements necessary to compute accurate solutions while increasing computational efficiency.

A MOVING MESH FINITE ELEMENT METHOD WITH LOCAL REFINEMENT FOR PARABOLIC PARTIAL DIFFERENTIAL EQUATIONS

Slimane Adjerd

**Department of Mathematical Science
Rensselaer Polytechnic Institute
Troy, New York 12180**

and

Joseph E. Flaherty

**Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180**

ABSTRACT

We discuss a moving mesh finite element method for solving initial-boundary value problems for vector systems of partial differential equations in one space dimension and time. The system is discretized using piecewise linear finite element approximations in space and a backward difference code for stiff ordinary differential systems in time. A spatial error estimation is calculated using piecewise quadratic approximations that use the superconvergence properties of parabolic systems to gain computational efficiency. The spatial error estimate is used to move and locally refine the finite element mesh in order to equidistribute a measure of the total spatial error and to satisfy a prescribed error tolerance. Ordinary differential equations for the spatial error estimate and the mesh motion are integrated in time using the same backward difference software that is used to determine the numerical solution of the partial differential system.

We present several details of an algorithm that may be used to develop a general purpose finite element code for one-dimensional parabolic partial differential systems. The algorithm combines mesh motion and local refinement in a relatively efficient manner and attempts to eliminate problem-dependent numerical parameters. A variety of examples that motivate our mesh moving strategy and illustrate the performance of our algorithm are presented.

**A DISCONTINUOUS FINITE ELEMENT
METHOD FOR HYPERBOLIC SYSTEMS
OF CONSERVATION LAWS**

T. L. Jackson

**Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180**

and

J. E. Flaherty

**Department of Computer Science
Rensselaer Polytechnic Institute
Troy, New York 12180**

ABSTRACT

We develop an adaptive finite element method for solving nonlinear hyperbolic systems of conservation laws in one space dimension and time. The method uses discontinuous piecewise linear trial functions and continuous piecewise cubic test functions on a moving mesh of triangular space-time elements. The mesh is moved by a technique that uses a weighted average of the local characteristic speeds to select nodal velocities. We show that discontinuous finite element approximations on a moving mesh have the potential of accurately resolving physical discontinuities in the solution. Several computational examples are presented to illustrate the performance of the method.

ON NONLINEAR SINGULARLY PERTURBED INITIAL VALUE PROBLEMS*

R. E. O'Malley, Jr.

Department of Mathematical Sciences
Rensselaer Polytechnic Institute
Troy, New York 12180

ABSTRACT

The study of singularly perturbed initial value problems for nonlinear systems of ordinary differential equations parallels the analysis underlying the development of numerical algorithms for obtaining solutions of such systems of stiff differential equations. This paper seeks to emphasize the advantages of combining these two substantial research efforts. It develops insight and intuition based on a sequence of solvable model problems, and it relates a variety of literature scattered throughout publications concerned with asymptotic and numerical analyses, stability and control theory, and specific topics in applied mathematical modeling.

*This material was presented as an invited talk at the Joint U.S.-Scandinavian Symposium on Scientific Computing and Mathematical Modeling in Stockholm, January 1985. The work was supported in part by the National Science Foundation under Grant Number DMS-8301665 and by the U. S. Army Research Office under Grant Number DAAG29-82-K-0197.

END

DTIC

7-86